



Unfavorable Structural and Functional Outcomes in Myelin Oligodendrocyte Glycoprotein Antibody-Associated Optic Neuritis

Jelcic, Ilijas ; Hanson, James V M ; Lukas, Sebastian ; Weber, Konrad P ; Landau, Klara ; Pless, Misha ; Reindl, Markus ; Weller, Michael ; Martin, Roland ; Lutterotti, Andreas ; Schippling, Sven

Abstract: BACKGROUND Recurrent optic neuritis (rON) associated with myelin oligodendrocyte glycoprotein (MOG)-specific antibodies has been initially reported to show a better clinical outcome than aquaporin-4 (AQP4)-seropositive ON in neuromyelitis optica spectrum disorder (NMOSD). Here, we characterize clinical and neuroimaging findings in severe cases of MOG antibody-positive and AQP4 antibody-negative bilateral rON. METHODS Three male adults with rON (ages 18, 44, and 63 years) were evaluated with optical coherence tomography (OCT), MRI, cerebrospinal fluid (CSF), and serological studies. RESULTS All patients experienced >7 relapses of ON with severe reduction of visual acuity and partial response to steroid treatment. Optic nerves were affected bilaterally, although unilateral relapses were more frequent than simultaneous bilateral recurrences. Patients were MOG-seropositive but repeatedly tested negative for AQP4 antibodies. OCT showed severe thinning of the peripapillary retinal nerve fiber layer. On MRI, contrast-enhancing lesions extended over more than half the length of the optic nerve. CSF analyses during ON episodes were normal. Severe visual deficits accumulated over time in 2 of 3 patients, despite immunosuppressive therapy. CONCLUSIONS MOG-seropositive and AQP4-seronegative rON may be associated with an aggressive disease course and poor functional and structural outcomes. In contrast to previous reports, the severity and pattern of retinal and optic nerve damage closely resembled phenotypes commonly observed in AQP4-seropositive rON without fulfilling current diagnostic criteria for NMOSD.

DOI: <https://doi.org/10.1097/WNO.0000000000000669>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-153699>

Journal Article

Published Version

Originally published at:

Jelcic, Ilijas; Hanson, James V M; Lukas, Sebastian; Weber, Konrad P; Landau, Klara; Pless, Misha; Reindl, Markus; Weller, Michael; Martin, Roland; Lutterotti, Andreas; Schippling, Sven (2019). Unfavorable Structural and Functional Outcomes in Myelin Oligodendrocyte Glycoprotein Antibody-Associated Optic Neuritis. *Journal of Neuro-Ophthalmology*, 39(1):3-7.

DOI: <https://doi.org/10.1097/WNO.0000000000000669>

Unfavorable Structural and Functional Outcomes in Myelin Oligodendrocyte Glycoprotein Antibody–Associated Optic Neuritis

Ilijas Jelcic, MD, James V. M. Hanson, PhD, Sebastian Lukas, Konrad P. Weber, MD, Klara Landau, MD, Misha Pless, MD, Markus Reindl, MD, Michael Weller, MD, Roland Martin, MD, Andreas Lutterotti, MD, Sven Schippling, MD

Background: Recurrent optic neuritis (rON) associated with myelin oligodendrocyte glycoprotein (MOG)-specific antibodies has been initially reported to show a better clinical outcome than aquaporin-4 (AQP4)-seropositive ON in neuromyelitis optica spectrum disorder (NMOSD). Here, we characterize clinical and neuroimaging findings in severe cases of MOG antibody–positive and AQP4 antibody–negative bilateral rON.

Methods: Three male adults with rON (ages 18, 44, and 63 years) were evaluated with optical coherence tomography (OCT), MRI, cerebrospinal fluid (CSF), and serological studies.

Results: All patients experienced >7 relapses of ON with severe reduction of visual acuity and partial response to steroid treatment. Optic nerves were affected bilaterally, although unilateral relapses were more frequent than simul-

taneous bilateral recurrences. Patients were MOG-seropositive but repeatedly tested negative for AQP4 antibodies. OCT showed severe thinning of the peripapillary retinal nerve fiber layer. On MRI, contrast-enhancing lesions extended over more than half the length of the optic nerve. CSF analyses during ON episodes were normal. Severe visual deficits accumulated over time in 2 of 3 patients, despite immunosuppressive therapy.

Conclusions: MOG-seropositive and AQP4-seronegative rON may be associated with an aggressive disease course and poor functional and structural outcomes. In contrast to previous reports, the severity and pattern of retinal and optic nerve damage closely resembled phenotypes commonly observed in AQP4-seropositive rON without fulfilling current diagnostic criteria for NMOSD.

Journal of Neuro-Ophthalmology 2018;0:1–5

doi: 10.1097/WNO.0000000000000669

© 2018 by North American Neuro-Ophthalmology Society

Neuroimmunology and Multiple Sclerosis Research (IJ, JVMH, SL, RM, AL, and SS), Neurology Clinic, University Hospital Zurich, University of Zurich, Zurich, Switzerland; Department of Ophthalmology (JVMH, KPW, and KL), University Hospital Zurich, Zurich, Switzerland; Department of Ophthalmology (MP), Luzerner Kantonsspital, Lucerne, Switzerland; Clinical Department of Neurology (MR), Medical University of Innsbruck, Innsbruck, Austria; and Neurology Clinic (MW, KPW), University Hospital Zurich, University of Zurich, Zurich, Switzerland.

I. Jelcic, J. V. M. Hanson, K. Landau, R. Martin, A. Lutterotti, and S. Schippling are supported by the Clinical Research Priority Project Multiple Sclerosis (CRPP^{MS}) of the University Zurich. S. Schippling and J. V. M. Hanson are supported by the Swiss MS Society. Before and subsequently to the conception and execution of the study, S. Lukas has been an employee of Heidelberg Engineering, Heidelberg, Germany.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.jneuro-ophthalmology.com).

A. Lutterotti and S. Schippling are equally contributing authors.

Address correspondence to Sven Schippling, MD, Neuroimmunology and Multiple Sclerosis Research Section, Department of Neurology, University Hospital Zurich, University of Zurich, Frauenklinikstrasse 26, 8091 Zurich, Switzerland; E-mail: Sven.schippling@usz.ch.

Myelin oligodendrocyte glycoprotein (MOG)-specific antibodies have been detected in peripheral blood of pediatric patients with acute disseminated encephalomyelitis (ADEM), but also in adults with benign, unilateral cerebral cortical encephalitis, aquaporin-4 (AQP4)-seronegative neuromyelitis optica spectrum disorder (NMOSD), AQP4-seronegative brainstem encephalitis, AQP4-seronegative (often longitudinally extensive) transverse myelitis, or in subjects with AQP4-seronegative optic neuritis (ON) (1–7). Because classic NMOSD is considered an astrocytopathy, and MOG antibody–associated diseases primarily display demyelination in the central nervous system, it has been a matter of debate whether MOG-seropositive, AQP4-seronegative syndromes should be classified as NMOSD, or whether they rather represent a separate autoimmune disease entity (8,9). MOG-seropositivity has been reported to be more frequently associated with simultaneous bilateral ON,

a monophasic disease course and a better functional outcome as compared to AQP4-seropositivity (10–14). Only recently have severe MOG antibody-associated recurrent ON (rON) been described (5,9,15,16). We evaluated 3 adult male patients with severe MOG antibody-positive rON and our aggressive immunosuppressive treatment attempts that could not prevent significant structural and/or functional sequelae.

METHODS

Patients provided written informed consent according to Swiss legislation; the use of clinical data was approved by the Cantonal Ethical Committee of Zurich, Switzerland (EC-ZH-No. 2013-0001). All patients received MRI and cerebrospinal fluid (CSF) examinations as routine diagnostic workup. Mean peripapillary retinal nerve fiber layer (pRNFL) thickness and macular volumes were assessed by optical coherence tomography (OCT) using a Spectralis SD-OCT device (Heidelberg Engineering, Heidelberg, Germany). Visual acuity (VA) was measured using Snellen charts; high- and low-contrast VA were assessed using ETDRS-style and 2.5% contrast SLOAN charts, respectively. Serum anti-MOG and anti-AQP4 antibodies were measured at the Clinical Department of Neurology, Innsbruck Medical University, by a cell-based assay as previously described (3). All patients were diagnosed with MOG-associated rON after excluding other diagnoses.

CASE REPORTS

Patient 1

A 18-year-old man experienced ON with subacute, painful visual loss in his right eye to hand movements. After high-dose corticosteroid, acuity in the right eye improved to 20/200. One month after the onset of ON, he experienced focal motor sensory and secondary generalized seizures. CSF analysis revealed normal cell count and protein levels without oligoclonal bands (OCBs). Brain MRI showed contrast enhancement over half the lengths of both optic nerves and the optic chiasm, as well as multiple T2- and contrast-enhancing lesions in the corpus callosum. Epileptic seizures subsided when the patient was treated with levetiracetam. Over the next 4 months, he experienced an episode of chiasmitis and 3 episodes of left ON. Treatments included high-dose corticosteroids and plasma exchange on 5 occasions.

Six months after his initial episodes of visual loss, he was referred to our institution. VA was 20/125, right eye, and worse than 20/400, left eye. pRNFL thickness was reduced in both eyes (right 46 μm and left 84 μm), with preferential thinning of the papillomacular bundle and the nasal and temporal quadrants (see **Supplemental Digital Content 2**, Fig. E1, <http://links.lww.com/WNO/A308>). Sera obtained on 3 occasions over a span of 10 months revealed MOG

antibodies with titers ranging from 1:160 to 1:640. AQP4-antibody tests were repeatedly negative. MOG antibody-associated chronic relapsing inflammatory optic neuropathy (CRION) was diagnosed.

Over the 6 months after our initial evaluation, the patient experienced 1 episode of right ON and 2 of left ON, despite continuous treatment with oral steroids (5–100 mg/day), use of rituximab $2 \times 375 \text{ mg/m}^2$, and plasma exchange. Notably, these episodes occurred during tapering the dose of steroids, and each responded moderately to high-dose oral steroids (100 mg/day over 5 days). Nine months after his initial episode, MRI showed persistent contrast enhancement of both optic nerves.

During the next 7 months, mycophenolate mofetil (up to 3,000 mg/day) was added as additional therapy. MRI no longer showed optic nerve enhancement. After repeating rituximab ($1 \times 375 \text{ mg/m}^2$), disease activity stabilized. At last follow-up, VA was 20/200, right eye, and no light perception, left eye, with pRNFL thickness of 28 μm in each eye.

In summary, 5 of 9 ON events were not only severe, but led to severe residual visual deficits. Rituximab therapy reduced relapse frequency, but could not prevent legal blindness (acuity worse than 20/200) as final outcome. The addition of mycophenolate mofetil to rituximab ameliorated the visual deficit on the right eye over time (see **Supplemental Digital Content 1**, Table E1, <http://links.lww.com/WNO/A307>).

Patient 2

A 43-year-old man was referred with an 8-year history of severe, painful rON. He initially experienced right ON with MRI showing edema and contrast enhancement along the entire right optic nerve. CSF analysis was normal. Subsequently, he had 1 additional episode of right ON, 4 of left ON, and 1 episode occurred bilaterally resulting in legal blindness. Although VA during ON episodes was reduced to as low as finger counting on both sides, the patient initially responded well to high-dose corticosteroid treatment. Therapy with azathioprine over 6 months was unsuccessful in preventing ON relapses. Repeat MRIs over 5 years showed exclusive optic nerve enhancement, whereas the spinal cord seemed normal. Multiple CSF analyses were normal. A diagnosis of CRION was made.

On our examination, VA was 20/20, right eye, and 20/32, left eye. OCT revealed pRNFL thinning (right: 48 μm ; left: 52 μm) (See **Supplemental Digital Content 2**, Fig. E1, <http://links.lww.com/WNO/A308>). The patient ultimately stabilized, after rituximab (375 mg/m^2) was given (See **Supplemental Digital Content 1**, Table E1, <http://links.lww.com/WNO/A307>). Most recently, acuity was 20/16 in both eyes. On 2 occasions, serum tested positive for MOG antibodies (titers 1:1,280 and 1:2,560) and negative for AQP4 antibodies.

Generally, Patient 2 recovered well after each of 10 ON events including severely disabling events. Rituximab

treatment reduced relapse frequency and may have stabilized the disease course in this patient.

Patient 3

A 59-year-old man experienced subacute back pain, urinary retention, saccadic smooth pursuit, and fever 6 weeks after an influenza vaccination. CSF analysis revealed an elevated cell count (183/ μ L) without OCB. MRI showed multiple T2-hyperintense bilateral thalamic, mesencephalic, pontine, and cervical, thoracic and lumbar spinal lesions with contrast enhancement of the thalamic, mesencephalic, pontine, and thoracic lesions. Microbiology and virology testing was negative, as was screening for AQP4 antibodies. A diagnosis of ADEM was made.

Two months later, he experienced painful right ON, with VA of 20/40. MRI demonstrated edema and contrast enhancement of almost the entire length of the right optic nerve. He was treated with high-dose corticosteroids, yet suffered a relapse of painful right ON. One month later, natalizumab (300 mg monthly) was added to the treatment regimen, but 3 further relapses of painful right ON occurred within 2 months. With a combination of intravenous high-dose steroids and plasma exchange, VA recovered to 20/40 in the right eye. Two months later, he received rituximab (375 mg/m²), and 1 month later, another 1000 mg.

Nevertheless, 2 months after the last rituximab infusion, he developed painful left ON and was given monthly infusions of cyclophosphamide (600–1100 mg/m²/cycle) for 13 months. During that time, the patient had 2 additional episodes of right ON. Repeat CSF analysis was unremarkable. Monthly infusions of tocilizumab (8 mg/kg) led to 17 months without relapse. VA was light perception, right eye, and 20/25, left eye. Measurement of pRNFL thickness was 30 μ m, right eye, with thinning over all quadrants (See **Supplemental Digital Content 2**, Fig E1, <http://links.lww.com/WNO/A308>), and 62 μ m, left eye, with predominant thinning over the nasal and inferior quadrants. MRI demonstrated atrophy of the right optic nerve and multiple supratentorial, T2-hyperintense lesions, but resolution of all spinal cord lesions. MOG antibodies were detected at a titer of 1:160 on 2 occasions but negative for AQP4 antibodies.

In summary, 4 of 8 ON events were severe and led to disabling residual visual deficits. Rituximab and cyclophosphamide treatment, respectively, reduced relapse frequency, but could not prevent severe ON relapses. Only initiation of tocilizumab therapy stabilized the disease completely without any further relapses or visual deterioration (See **Supplemental Digital Content 1**, Table E1, <http://links.lww.com/WNO/A307>).

DISCUSSION

We present 3 adult male patients with MOG antibody-associated rON (MOG-ON) with poor structural out-

comes in all cases and poor functional outcomes in 2 of 3 cases. None of our patients fulfilled current diagnostic criteria for NMOSD (17). Two patients experienced an ADEM-like episode after (Patient 1) or before (Patient 3) their first episode of ON. In accordance with previous reports (16,18), CSF findings were normal in 2 patients during MOG-ON episodes. The initial elevated CSF cell count and mild blood–brain barrier dysfunction in Patient 3 may be related to the antecedent influenza vaccination and suggestive of both MOG-seropositive ADEM and NMOSD, as both have been reported to feature increased CSF cell counts (12,19). To the best of our knowledge, this is only the third case with vaccination-related MOG-seropositive disease (9). Other MOG-seropositive cases have been associated with viral or bacterial infection including influenza virus infection, Epstein–Barr virus-induced infectious mononucleosis and *Chlamydia pneumoniae* infection (20–22). In these cases, autoreactive MOG antibody-producing B cells may be activated by both the simultaneous uptake of the cognate autoantigen (MOG) and the “bystander” viral antigen (e.g., influenza hemagglutinin) into B cells from infected parenchymal cells and by T cells specific for the respective antigen (23).

Extensive contrast enhancement of the optic nerve (frequently extending over more than half of the entire length and involving the chiasm) on MRI and marked thinning of the entire pRNFL resembled typical findings in NMOSD (24). OCT studies have shown preferential temporal pRNFL thinning in multiple sclerosis-associated rON, whereas the initial involvement of the superior and inferior quadrants is typical in NMOSD-associated rON along with severe macular thinning (24,25). By contrast, in all 3 patients with MOG-ON reported here, pRNFL thickness was reduced in all quadrants.

Tapering of corticosteroid therapy led to ON relapses in Patient 1. Dependency on long-term corticosteroid treatment and good responsiveness to high-dose corticosteroid treatment are typical of CRION. MOG-seropositive CRION and rON cases with rapid steroid response and increased relapse risk after steroid cessation have been reported (4,9,16). Notably, Patients 1 and 2 showed only partial clinical response to antibody-targeted therapy with rituximab.

Within the past few years, the first MOG-ON cases with the onset of >18 years of age have been reported (15,16). These patients tended to show worse VA outcome than pediatric cases (15). However, only 2 of 14 patients in these reports (Case 10 (15); Case 7 (16)) experienced such a severe reduction of VA like the cases presented here. Subsequently, other cases with recurrent (14,27) and/or severe disease course have been described (9,28,29). As reports of MOG-seropositive patients with longer follow-up (>5 years) become available, poor visual outcomes will likely become more prevalent.

Because MOG-ON reportedly affects female patients more often than male patients, with higher annualized relapse rate in female patients (9), the cases reported here (all men and frequent rON) may represent one end of the spectrum of MOG-ON. Whether the later onset of MOG-ON and/or gender of the individual is associated with a more aggressive disease course warrants further study. Testing for MOG antibodies should be considered in cases of rON or suspected NMOSD, particularly in patients with AQP4-seronegative. Notably, MOG antibody titers did not relate to ON severity or pRNFL loss. Future studies including larger case series will have to address the pathological relevance and overall clinical utility of MOG-antibody testing for clinical decision making.

In summary, our cases suggest that MOG-ON may be associated with disabling structural and functional visual outcomes, similar to patients with AQP4-seropositive ON. Initiation of high efficacy immune therapy should be evaluated early in these patients, especially when initial episodes are followed by poor visual outcome.

STATEMENT OF AUTHORSHIP

Category 1: a. Conception and design: I. Jelcic, J. V. M. Hanson, S. Lukas, A. Lutterotti, and S. Schippling; b. Acquisition of data: I. Jelcic, J. V. M. Hanson, S. Lukas, K. P. Weber, M. Pless, M. Reindl, A. Lutterotti, and S. Schippling; c. Analysis and interpretation of data: I. Jelcic, J. V. M. Hanson, S. Lukas, K. P. Weber, K. Landau, M. Pless, M. Reindl, M. Weller, R. Martin, A. Lutterotti, and S. Schippling; Category 2: a. Drafting the manuscript: I. Jelcic, J. V. M. Hanson, A. Lutterotti, and S. Schippling; b. Revising it for intellectual content: I. Jelcic, J. V. M. Hanson, S. Lukas, K. P. Weber, K. Landau, M. Pless, M. Reindl, M. Weller, R. Martin, A. Lutterotti, and S. Schippling; Category 3: a. Final approval of the completed manuscript: I. Jelcic, J. V. M. Hanson, S. Lukas, K. P. Weber, K. Landau, M. Pless, M. Reindl, M. Weller, R. Martin, A. Lutterotti, and S. Schippling.

ACKNOWLEDGMENTS

The authors thank Kathrin Schanda, Clinical Department of Neurology, Medical University of Innsbruck, Innsbruck, Austria, for excellent technical assistance.

REFERENCES

1. **Brilot F**, Dale RC, Selzer RC, Grummel V, Kalluri SR, Aslam M, Busch V, Zhou D, Cepok S, Hemmer B. Antibodies to native myelin oligodendrocyte glycoprotein in children with inflammatory demyelinating central nervous system disease. *Ann Neurol*. 2009;66:833–842.
2. **Kitley J**, Woodhall M, Waters P, Leite MI, Devenney E, Craig J, Palace J, Vincent A. Myelin-oligodendrocyte glycoprotein antibodies in adults with a neuromyelitis optica phenotype. *Neurology*. 2012;79:1273–1277.
3. **Mader S**, Gredler V, Schanda K, Rostasy K, Dujmovic I, Pfaller K, Lutterotti A, Jarius S, Di Pauli F, Kuenz B, Ehling R, Hegen H, Deisenhammer F, Aboul-Enein F, Storch MK, Koson P, Drulovic J, Kristoferitsch W, Berger T, Reindl M. Complement activating antibodies to myelin oligodendrocyte glycoprotein in neuromyelitis optica and related disorders. *J Neuroinflammation*. 2011;8:184.

4. **Mayer MC**, Breithaupt C, Reindl M, Schanda K, Rostasy K, Berger T, Dale RC, Brilot F, Olsson T, Jenne D, Probstel AK, Dornmair K, Wekerle H, Hohlfeld R, Banwell B, Bar-Or A, Meinl E. Distinction and temporal stability of conformational epitopes on myelin oligodendrocyte glycoprotein recognized by patients with different inflammatory central nervous system diseases. *J Immunol*. 2013;191:3594–3604.
5. **Rostasy K**, Mader S, Schanda K, Huppke P, Gartner J, Kraus V, Karenfort M, Tibussek D, Blaschek A, Bajer-Kornek B, Leitz S, Schimmel M, Di Pauli F, Berger T, Reindl M. Anti-myelin oligodendrocyte glycoprotein antibodies in pediatric patients with optic neuritis. *Arch Neurol*. 2012;69:752–756.
6. **Ogawa R**, Nakashima I, Takahashi T, Kaneko K, Akaishi T, Takai Y, Sato DK, Nishiyama S, Misu T, Kuroda H, Aoki M, Fujihara K. MOG antibody-positive, benign, unilateral, cerebral cortical encephalitis with epilepsy. *Neurol Neuroimmunol Neuroinflamm*. 2017;4:e322.
7. **Jarius S**, Kleiter I, Ruprecht K, Asgari N, Pitarokoli K, Borisow N, Hummert MW, Trebst C, Pache F, Winkelmann A, Beume LA, Ringelstein M, Stich O, Aktas O, Korpall-Kuhnke M, Schwarz A, Lukas C, Haas J, Fechner K, Buttman M, Bellmann-Strobl J, Zimmermann H, Brandt AU, Franciotta D, Schanda K, Paul F, Reindl M, Wildemann B; in Cooperation With the Neuromyelitis Optica Study G. MOG-IgG in NMO and related disorders. A multicenter study of 50 patients. Part 3: brainstem involvement—frequency, presentation and outcome. *J Neuroinflammation*. 2016;13:281.
8. **Zamvil SS**, Slavin AJ. Does MOG Ig-positive AQP4-seronegative opticospinal inflammatory disease justify a diagnosis of NMO spectrum disorder? *Neurol Neuroimmunol Neuroinflamm*. 2015;2:e62.
9. **Jarius S**, Ruprecht K, Kleiter I, Borisow N, Asgari N, Pitarokoli K, Pache F, Stich O, Beume LA, Hummert MW, Ringelstein M, Trebst C, Winkelmann A, Schwarz A, Buttman M, Zimmermann H, Kuchling J, Franciotta D, Capobianco M, Siebert E, Lukas C, Korpall-Kuhnke M, Haas J, Fechner K, Brandt AU, Schanda K, Aktas O, Paul F, Reindl M, Wildemann B; in Cooperation With the Neuromyelitis Optica Study G. MOG-IgG in NMO and related disorders. A multicenter study of 50 patients. Part 2: epidemiology, clinical presentation, radiological and laboratory features, treatment responses, and long-term outcome. *J Neuroinflammation*. 2016;13:280.
10. **Hoftberger R**, Sepulveda M, Armangue T, Blanco Y, Rostasy K, Cobo Calvo A, Olascoaga J, Ramio-Torrenta L, Reindl M, Benito-Leon J, Casanova B, Arrambide G, Sabater L, Graus F, Dalmau J, Saiz A. Antibodies to MOG and AQP4 in adults with neuromyelitis optica and suspected limited forms of the disease. *Mult Scler*. 2015;21:866–874.
11. **Kitley J**, Waters P, Woodhall M, Leite MI, Murchison A, George J, Kuker W, Chandratte S, Vincent A, Palace J. Neuromyelitis optica spectrum disorders with aquaporin-4 and myelin-oligodendrocyte glycoprotein antibodies: a comparative study. *JAMA Neurol*. 2014;71:276–283.
12. **Sato DK**, Callegaro D, Lana-Peixoto MA, Waters PJ, de Haidar Jorge FM, Takahashi T, Nakashima I, Apostolos-Pereira SL, Talim N, Simm RF, Lino AM, Misu T, Leite MI, Aoki M, Fujihara K. Distinction between MOG antibody-positive and AQP4 antibody-positive NMO spectrum disorders. *Neurology*. 2014;82:474–481.
13. **Stiebel-Kalish H**, Lotan I, Brody J, Chodick G, Bialer O, Marignier R, Bach M, Hellmann MA. Retinal nerve fiber layer may be better preserved in MOG-IgG versus AQP4-IgG optic neuritis: a cohort study. *PLoS One*. 2017;12:e0170847.
14. **Akaishi T**, Sato DK, Nakashima I, Takeshita T, Takahashi T, Doi H, Kurosawa K, Kaneko K, Kuroda H, Nishiyama S, Misu T, Nakazawa T, Fujihara K, Aoki M. MRI and retinal abnormalities in isolated optic neuritis with myelin oligodendrocyte glycoprotein and aquaporin-4 antibodies: a comparative study. *J Neurol Neurosurg Psychiatry*. 2016;87:446–448.
15. **Martinez-Hernandez E**, Sepulveda M, Rostasy K, Hoftberger R, Graus F, Harvey RJ, Saiz A, Dalmau J. Antibodies to aquaporin 4, myelin-oligodendrocyte glycoprotein, and the glycine

- receptor alpha1 subunit in patients with isolated optic neuritis. *JAMA Neurol.* 2015;72:187–193.
16. **Ramanathan S**, Reddel SW, Henderson A, Parratt JD, Barnett M, Gatt PN, Merheb V, Kumaran RY, Pathmanandavel K, Sinmaz N, Ghadiri M, Yiannikas C, Vucic S, Stewart G, Bleasel AF, Booth D, Fung VS, Dale RC, Brilot F. Antibodies to myelin oligodendrocyte glycoprotein in bilateral and recurrent optic neuritis. *Neurol Neuroimmunol Neuroinflamm.* 2014;1:e40.
 17. **Wingerchuk DM**, Banwell B, Bennett JL, Cabre P, Carroll W, Chitnis T, de Seze J, Fujihara K, Greenberg B, Jacob A, Jarius S, Lana-Peixoto M, Levy M, Simon JH, Tenenbaum S, Traboulsee AL, Waters P, Wellik KE, Weinshenker BG; International Panel for NMOS. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology.* 2015;85:177–189.
 18. **Nakajima H**, Motomura M, Tanaka K, Fujikawa A, Nakata R, Maeda Y, Shima T, Mukaino A, Yoshimura S, Miyazaki T, Shiraishi H, Kawakami A, Tsujino A. Antibodies to myelin oligodendrocyte glycoprotein in idiopathic optic neuritis. *BMJ Open.* 2015;5:e007766.
 19. **Kortvelyessy P**, Breu M, Pawlitzki M, Metz I, Heinze HJ, Matzke M, Mawrin C, Rommer P, Kovacs GG, Mitter C, Reindl M, Bruck W, Wandinger KP, Lassmann H, Hoftberger R, Leypoldt F. ADEM-like presentation, anti-MOG antibodies, and MS pathology: two case reports. *Neurol Neuroimmunol Neuroinflamm.* 2017;4:e335.
 20. **Amano H**, Miyamoto N, Shimura H, Sato DK, Fujihara K, Ueno S, Nakamura R, Ueno Y, Watanabe M, Hattori N, Urabe T. Influenza-associated MOG antibody-positive longitudinally extensive transverse myelitis: a case report. *BMC Neurol.* 2014;14:224.
 21. **Nakamura Y**, Nakajima H, Tani H, Hosokawa T, Ishida S, Kimura F, Kaneko K, Takahashi T, Nakashima I. Anti-MOG antibody-positive ADEM following infectious mononucleosis due to a primary EBV infection: a case report. *BMC Neurol.* 2017;17:76.
 22. **Azumagawa K**, Nomura S, Shigeri Y, Jones LS, Sato DK, Nakashima I, Kashiwagi M, Tanabe T, Shimakawa S, Nakajima H, Tamai H. Post-vaccination MDEM associated with MOG antibody in a subclinical Chlamydia infected boy. *Brain Dev.* 2016;38:690–693.
 23. **Sanderson NS**, Zimmermann M, Eilinger L, Gubser C, Schaeren-Wiemers N, Lindberg RL, Dougan SK, Ploegh HL, Kappos L, Derfuss T. Cocapture of cognate and bystander antigens can activate autoreactive B cells. *Proc Natl Acad Sci U S A.* 2017;114:734–739.
 24. **Bennett JL**, de Seze J, Lana-Peixoto M, Palace J, Waldman A, Schippling S, Tenenbaum S, Banwell B, Greenberg B, Levy M, Fujihara K, Chan KH, Kim HJ, Asgari N, Sato DK, Saiz A, Wuerfel J, Zimmermann H, Green A, Villoslada P, Paul F; Gjc ICC, Br. Neuromyelitis optica and multiple sclerosis: seeing differences through optical coherence tomography. *Mult Scler.* 2015;21:678–688.
 25. **Schneider E**, Zimmermann H, Oberwahrenbrock T, Kaufhold F, Kadas EM, Petzold A, Bilger F, Borisow N, Jarius S, Wildemann B, Ruprecht K, Brandt AU, Paul F. Optical coherence tomography reveals distinct patterns of retinal damage in neuromyelitis optica and multiple sclerosis. *PLoS One.* 2013;8:e66151.
 26. **Huppke P**, Rostasy K, Karenfort M, Huppke B, Seidl R, Leiz S, Reindl M, Gartner J. Acute disseminated encephalomyelitis followed by recurrent or monophasic optic neuritis in pediatric patients. *Mult Scler.* 2013;19:941–946.
 27. **Chang T**, Waters P, Woodhall M, Vincent A. Recurrent optic neuritis associated with MOG antibody seropositivity. *Neurologist.* 2017;22:101–102.
 28. **Havla J**, Kumpfel T, Schinner R, Spadaro M, Schuh E, Meinl E, Hohlfeld R, Outteryck O. Myelin-oligodendrocyte-glycoprotein (MOG) autoantibodies as potential markers of severe optic neuritis and subclinical retinal axonal degeneration. *J Neurol.* 2017;264:139–151.
 29. **Pache F**, Zimmermann H, Mikolajczak J, Schumacher S, Lacheta A, Oertel FC, Bellmann-Strobl J, Jarius S, Wildemann B, Reindl M, Waldman A, Soelberg K, Asgari N, Ringelstein M, Aktas O, Gross N, Buttmann M, Ach T, Ruprecht K, Paul F, Brandt AU; in Cooperation With the Neuromyelitis Optica Study G. MOG-IgG in NMO and related disorders. A multicenter study of 50 patients. Part 4: afferent visual system damage after optic neuritis in MOG-IgG-seropositive versus AQP4-IgG-seropositive patients. *J Neuroinflammation.* 2016;13:282.